

MAINE MONTHLY OVERDOSE REPORT

For March 2026

Marcella H. Sorg, Prianka Maria Sarker, and Daniel S. Soucier
Margaret Chase Smith Policy Center, University of Maine


Overview

This report documents suspected and confirmed fatal and nonfatal drug overdoses in Maine for the period January 2025–March 2026 (Table 1). The total number of confirmed and suspected fatal overdoses for January–March 2026 is 83, 13.5% lower than the total confirmed fatal overdoses for January–March 2025, 96. The total number of nonfatal overdoses for January–March 2026 is 1,636, 2.6% lower compared to the reported nonfatal overdoses for January–March 2025, 1,679. Monthly proportions of fatalities during January–March 2026 averaged 4.8%. During 2025, monthly proportions of fatalities averaged 5.1%, ranging from a high of 6.5% in December to a low of 3.3% in September.

Data derived from multiple statewide sources were compiled and deduplicated to calculate fatal and nonfatal overdose totals (Table 1). These include nonfatal overdose incidents reported by hospital emergency departments (ED), nonfatal emergency medical service (EMS) responses without transport to the ED, overdose reversals reported by law enforcement in the absence of EMS, and overdose reversals reported by community members or agencies receiving state-supplied naloxone through the Maine Naloxone Distribution Initiative. There are also an unknown number of private overdose reversals that were not reported and an unknown number of community-reported reversals that may have overlapped with emergency response by EMS or law enforcement. The total number of fatal overdoses in this report includes those that have been confirmed, as well as those that are suspected but not yet confirmed for December 2025, February 2026, and March 2026 (see Figure 1).¹

The total number of suspected and confirmed fatal overdoses and reported nonfatal overdoses for March 2026, 607, is displayed in Table 1. Of those 607, there were 24 (4.0%) confirmed and suspected fatal overdoses, 302 (49.8%) nonfatal emergency department visits, 167 (27.5%) nonfatal EMS responses not transported to the

Figure 1. Suspected and confirmed fatal overdoses, all drugs, January 2025 through March 2026



¹ Until overdose is confirmed as a cause of death, the manner of death is not certified. The vast majority of the suspected fatal overdoses are ultimately certified accidental manner of death, approximately 90%.

emergency department, 97 (16.0%) reported community overdose reversals, and 17 (2.8%) law enforcement reversals in incidents that did not include EMS.

Note that we have modified reporting methods for subpopulations. Data tables now report a rolling three-month total rather than a total for a single month. This improves confidentiality protections and reduces the focus on very small totals that are not statistically significant. In addition, we are now suppressing subpopulation totals less than 5.

Table 1: Composite reported overdose totals, all drugs, January 2025–March 2026

	Nonfatal					Total confirmed and suspected fatal overdoses	Total overdoses
	Emergency Dept.	EMS not transported to emergency	Community reversals with naloxone	Law enforcement nonfatal overdose response without EMS	Total nonfatal overdoses		
January 2025	234	193	53	15	495	31	526
February 2025	254	159	74	18	505	33	538
March 2025	375	202	71	31	679	32	711
April 2025	295	212	90	18	615	36	651
May 2025	311	202	88	13	614	30	644
June 2025	399	233	96	21	749	40	789
July 2025	321	247	70	15	653	32	685
August 2025	272	191	76	13	552	31	583
September 2025	385	244	118	13	760	26	786
October 2025	254	194	100	11	559	28	587
November 2025	267	177	70	9	523	33	556
December 2025	319	178	104	8	609	42	651
2025 YTD Total	3686	2432	1010	185	7313	394	7707
% of 2025 YTD Total	47.8%	31.6%	13.1%	2.4%	94.9%	5.1%	100%
January 2026	257	156	105	14	532	28	560
February 2026	264	149	96	12	521	31	552
March 2026	302	167	97	17	583	24	607
2026 YTD Total	823	472	298	43	1636	83	1719
% of 2026 YTD Total	47.9%	27.5%	17.3%	2.5%	95.2%	4.8%	100%

Law Enforcement Response to Fatal and Nonfatal Overdose Incidents

Due to the method used to deduplicate nonfatal overdose incidents to derive a composite number of overdoses for the month, the total activity of both law enforcement officials and EMS agencies is underrepresented in Table 1 because of the overlap between them. The process used to deduplicate overdoses begins by removing fatal overdoses from the emergency department and EMS overdose incidents. Then the number of patients transported to emergency departments by Maine EMS are removed from the EMS overdose incidents. Finally, EMS involvement and fatal overdose incidents are removed from law enforcement responses.

Table 2 shows the public safety response to fatal and nonfatal overdose events in January–March 2026 as well as during January–December 2025. During 2025, law enforcement officers responded to a reported 862 overdose incidents (328 fatal; 534 nonfatal), and Maine EMS responded to a reported 9,172 incidents (297 fatal; 8,875 nonfatal). In January–March 2026, law enforcement officers responded to a reported 169 overdose incidents (71 fatal; 98 nonfatal), and Maine EMS responded to a reported 1,951 incidents (57 fatal; 1,894 nonfatal).

Table 2: Fatal and nonfatal overdose emergency response counts from law enforcement and EMS, including overlapping cases

	Fatal overdose response Jan-Dec 2025	Nonfatal overdose response Jan-Dec 2025	Total overdose response Jan-Dec 2025	Fatal overdose response Jan-Mar 2026	Nonfatal overdose response Jan-Mar 2026	Total overdose response Jan-Mar 2026
Maine EMS	297	8875	9172	57	1894	1951
Monthly average	25	740	764	19	631	650
Law Enforcement	328	534	862	71	98	169
Monthly average	27	45	72	24	33	56

Note: Numbers will fluctuate from month to month as public safety agencies catch up their reporting. Due to methodological convention, alcohol-only cases are excluded from this table. However, we recognize that alcohol is a large part of substance misuse epidemic. Cases with both drugs and alcohol are included. Please note these numbers may fluctuate higher than the data in Table 1. This is due to the fact that some EMS overdoses responses, once the patient is transported to the hospital, are deemed to be non-overdose emergencies such as cardiac arrest or diabetic coma.

County Distribution of Suspected Nonfatal Overdoses with EMS Response

Table 3 shows the frequency distribution of nonfatal overdoses to which EMS responded at the county level. Overdose reversal totals reported by community partners and emergency departments are not reported by county; only EMS case data include county frequencies. To improve statistical reliability and protect data confidentiality, the county-level totals are presented as a yearly total for 2025 and as a rolling 3-month aggregate for the period January–March 2026. The rolling 3-month aggregate percentage in the rightmost column can be compared to the percentage of the census population, or to the percentage of nonfatal overdoses for the year 2025. Caution must be exercised viewing single counties, even the three-month totals, as these totals may fluctuate randomly, without reflecting any statistically significant trend.

During January–March 2026, percentages for most counties fall within 0 to 2 percentage points of the 2020 census distribution. Compared to the 2020 census proportion, Cumberland County is 5 percentage points higher, and Androscoggin County is 4 percentage points higher. York County is 3 percentage points lower.

Table 3: County of EMS incident among suspected and confirmed nonfatal overdoses

	% 2020 estimated Census population	Jan-Dec 2025 Est. N = 8875	Jan-Mar 2026 Est. N = 1893
Androscoggin	8%	1004 11%	228 12%
Aroostook	5%	372 4%	80 4%
Cumberland	22%	2374 27%	519 27%
Franklin	2%	152 2%	32 2%
Hancock	4%	236 3%	57 3%
Kennebec	9%	799 9%	152 8%
Knox	3%	257 3%	47 2%
Lincoln	3%	165 2%	40 2%
Oxford	4%	331 4%	71 4%
Penobscot	11%	1146 13%	230 12%
Piscataquis	1%	96 1%	19 1%
Sagadahoc	3%	144 2%	31 2%
Somerset	4%	358 4%	66 3%
Waldo	3%	194 2%	37 2%
Washington	2%	217 2%	39 2%
York	16%	1030 12%	245 13%

Note: EMS nonfatal overdose counts include incidents where a patient may have died after admission to the ED. Please note numbers will fluctuate from month-to-month as public safety agencies catch up their reporting. Due to methodological convention, alcohol-only cases are excluded from this table. However, we recognize that alcohol is a large part of substance misuse epidemic. Cases with both drugs and alcohol are included.

Age and Gender Distribution of Suspected Nonfatal Overdoses with EMS Response

Table 4 displays the age composition of individuals suspected of experiencing nonfatal overdoses involving EMS response. Overdose reversal totals reported by community partners and emergency departments are not categorized and reported by age; only EMS case data include age frequencies. To support statistical reliability and data confidentiality, subpopulation age group proportions are presented as annual figures for the year 2025 and as rolling 3-month totals for January–March 2026.

The age distribution for both the 2025 yearly total and the rolling 3-month total (January–March 2026) compared to the 2020 census proportion shows a disproportionately large impact of suspected nonfatal overdose victims with EMS involvement for those aged 25–54. This impact is illustrated by looking at the 25–54-year age groups, which constitute 36% of the population in the 2020 census compared to 62% of the nonfatal overdose population in 2025, and 60% of the nonfatal overdose population during January–March 2026. In the most recent three months, there are 14 percentage points fewer overdose victims among those under the age of 18 compared to the percentage of the census population in that age group. Similarly, there were 2 percentage points fewer overdose victims among those aged 55–64, and 9 percentage points fewer overdose victims among those 65 and older compared to the percentages of the census population for those age groups.

Table 4: Reported age group among suspected nonfatal overdose victims involving EMS response

	% 2020 estimated Census population	Jan-Dec 2025 Est. N = 8885		Jan- Mar 2026 Est. N = 1896	
< 18	18%	367	4%	76	4%
18-24	7%	726	8%	169	9%
25-34	12%	1585	18%	318	17%
35-44	12%	2423	27%	523	28%
45-54	12%	1531	17%	296	16%
55-64	16%	1091	12%	257	14%
> 64	23%	1162	13%	257	14%

Table 5 displays the reported gender of individuals experiencing nonfatal overdoses involving EMS response. Overdose reversal totals reported by community partners and emergency departments are not categorized by gender; only EMS case data include gender categories. To support statistical reliability and maintain data confidentiality, gender group proportions are presented as annual figures for the year 2025 and as rolling 3-month totals for January–March 2026.

Gender group proportions in the EMS data for the year 2025 or for the rolling 3-month aggregates can be compared to the 2020 census proportion in the leftmost column. Males represent 49% of the 2020 estimated census population and 59% of the nonfatal overdose victims with EMS involvement both in 2025 and in January–March 2026.

Table 5: Reported gender among suspected nonfatal overdose victims involving EMS response

	% 2020 estimated Census population	Jan-Dec 2025 Est. N = 8567		Jan-Mar 2026 Est. N = 1827	
Male	49%	5076	59%	1085	59%
Female	51%	3491	41%	742	41%

County and City Distribution of Suspected and Confirmed Fatal Overdoses

Table 6 shows the frequency distribution of fatal overdoses at the county level including larger metro cities.² To mitigate monthly volatility and confidentiality concerns due to small cell sizes, county/city level estimates are presented as yearly totals for 2025 and as rolling 3-month totals for January–March 2026, instead of monthly figures, and totals below 5 are suppressed.

During January–March 2026, the percentages for most counties fall within 0 to 3 percentage points of the 2020 census distribution. Compared to the 2020 census proportion, Cumberland County is 6 percentage points higher, Penobscot County is 3 percentage points higher, Washington County is 5 percentage points higher, and York County is 5 percentage points lower. At the city level for January–March 2026, Bangor is 6 percentage points and Portland is 14 percentage points higher than the 2020 census population.

² This table reports the county/city of death rather than the county/city of residence.

Table 6: County/city of death among suspected and confirmed fatal overdoses

	% 2020 estimated Census population	Jan-Dec 2025 Est. N = 394		Jan-Mar 2026 Est. N = 83	
Androscoggin	8%	41	10%	5	6%
Lewiston	3%	28	7%	<5	*
Aroostook	5%	27	7%	<5	*
Cumberland	22%	86	22%	23	28%
Portland	5%	51	13%	16	19%
Franklin	2%	7	2%	<5	*
Hancock	4%	6	2%	<5	*
Kennebec	9%	27	7%	6	7%
Knox	3%	9	2%	<5	*
Lincoln	3%	8	2%	<5	*
Oxford	4%	16	4%	<5	*
Penobscot	11%	69	18%	12	14%
Bangor	2%	36	9%	7	8%
Piscataquis	1%	<5	*	<5	*
Sagadahoc	3%	<5	*	<5	*
Somerset	4%	10	3%	6	7%
Waldo	3%	14	4%	<5	*
Washington	2%	14	4%	6	7%
York	16%	54	14%	9	11%

<5 Estimates with a count of less than 5 are suppressed due to statistical reliability and confidentiality purposes.

* Percentages are suppressed due to small cell sizes (n <5).

Age and Sex Distribution of Fatal Overdose Victims

Table 7 displays the age and sex composition³ of the fatal overdose population for the year 2025 and for the rolling 3-month period of January–March 2026.

The rolling 3-month aggregate proportion of males for January to March 2026 is 60%, which is 11 percentage points higher than the census population. The cumulative proportion of males in 2025 is 66%. The age distribution for both the year 2025 and the period January–March 2026 compared to the 2020 census proportion shows a disproportionately large impact of fatal overdoses in those aged 35–64. That group includes 40% of the 2020 estimated census population, compared to 72% of the fatal overdose population during January–March 2026 and 75% of the fatal overdose population in 2025. Those ages 18–24 constitute 7% of the 2020 census population but only 2% of 2025 fatal overdoses. Those 65 and older constitute 23% of the census population but only 11% of the 2025 fatal overdoses and 14% of the January–March 2026 fatal overdoses.

3 Note that death certificate reports sex as male or female without gender categories.

Table 7: Decedent reported age group and sex among suspected and confirmed fatal overdoses

	% 2020 estimated Census population	Jan-Dec 2025 Est. N = 394	Jan-Mar 2026 Est. N = 83
Male	49%	260 66%	50 60%
< 18	18%	<5 *	<5 *
18-24	7%	6 2%	<5 *
25-34	12%	49 12%	8 10%
35-44	12%	115 29%	28 34%
45-54	12%	98 25%	19 23%
55-64	16%	82 21%	13 16%
> 64	23%	44 11%	12 14%

<5 Estimates with a count of less than 5 are suppressed due to statistical reliability and confidentiality purposes.
 * Percentages are suppressed due to small cell sizes (n <5).
 Note: Percentages may not total 100 due to rounding.

Table 8 displays the reported race and ethnicity of confirmed and suspected fatal overdoses during January–December 2025 and January–March 2026, compared to the 2020 estimated census population. Note that race and ethnicity are not finalized until the full death certificate is entered into Vital Records, and a small number of decedents’ records currently lack information about these variables. Out of 83 decedents for whom race was reported in January–March 2026, 90% of the victims were identified as White.

Table 8: Decedent race and ethnicity among suspected and confirmed fatal overdoses⁴

A: Race	% 2020 estimated Census population	Jan-Dec 2025 Race N = 394	Jan-Mar 2026 Race N = 83
White alone	94%	379 96%	75 90%
Black/African American alone	2%	7 2%	<5 *
American Indian/Alaska Native alone	1%	<5 *	<5 *
Other race and 2+ races combined	3%	5 1%	<5 *
B: Ethnicity	% 2020 estimated Census population	Jan-Dec 2025 Ethnicity N = 391	Jan-Mar 2026 Ethnicity N = 82
Hispanic/Latinx alone	2%	<5 *	<5 *

<5 Estimates with a count of less than 5 are suppressed due to statistical reliability and confidentiality purposes.
 * Percentages are suppressed due to small cell sizes (n <5).
 Note: Percentages may not total 100 due to rounding.

Military Status and Housing Stability of Fatal Overdose Victims

Out of the 394 cases for which military background was reported in January–December 2025, 29 (7%) were identified as having a military background. Out of the 83 cases in January–March 2026 where military background was reported, fewer than 5 cases were identified as having a military background.

Of the 394 total suspected and confirmed fatal overdose cases in January–December 2025, undomiciled or transient housing status was reported for 50 (13%) victims. Of the 83 total suspected and confirmed fatal overdose cases in January–March 2026, fewer than 5 victims were identified as undomiciled.

⁴ Race and ethnicity data for some cases are unavailable until drug deaths are confirmed.

Basic Incident Patterns of Fatal Overdoses

Table 9 reports basic incident patterns for fatal overdoses. For statistical relevance and confidentiality purposes, totals are reported for 2025 as a whole and as rolling 3-month totals for January–March 2026. Caution must be exercised since data totals may change slightly as suspected cases are confirmed or eliminated.

Both EMS and police responded together to most fatal overdoses (55%) in January–March 2026. Law enforcement was more likely to respond to a scene alone (30%) than EMS (13%) was. Most (91%) suspected and confirmed fatal drug overdoses were accidental manner of death. Of the 77 confirmed or suspected fatal overdoses in January–March 2026 for which overdose history was known, 16 (21%) had a reported history of prior overdose.

In most overdose cases, bystanders or witnesses were present at the scene by the time first responders arrived, although the details about who was present at the time of the overdose were frequently unclear. In some cases, responding family and friends or other bystanders administered naloxone. Often, EMS and/or law enforcement administered naloxone in addition to bystanders or witnesses. During January–March 2026, 14% of the suspected and confirmed fatal overdose cases had naloxone administered at the scene by EMS, law enforcement, or bystanders, or by combinations of these entities.

Of the 57 suspected or confirmed drug death cases with EMS response during January–March 2026, 24 (42%) victims were already deceased when EMS arrived. In the remaining 33 (58%) cases, resuscitation was attempted either at the scene or presumably in the ambulance during transport to the emergency room. Of those 33 ultimately fatal victims who were alive when EMS arrived, 14 (42%) were transported, and 19 (58%) did not survive to be transported. Thus, out of 57 ultimately fatal cases with EMS response, only 14 (25%) remained alive long enough to be transported but died during transport or at the emergency room. This outcome is likely due to a combination of the high number of cases with fentanyl as a cause of death and individuals using alone. Fentanyl acts more quickly than other opioids, and there is less time for bystanders to find an overdose victim alive, administer naloxone, and call 911.

Table 9: Incident characteristics among suspected and confirmed fatal overdoses

	Jan–Dec 2025 Est. N = 394	Jan–Mar 2026 Est. N = 83
EMS response alone	57 14%	11 13%
Law enforcement alone	87 22%	25 30%
EMS and law enforcement	240 61%	46 55%
Naloxone administration reported at the scene	95 24%	12 14%
EMS administered naloxone	62 16%	<5 *
Law enforcement administered naloxone	22 6%	<5 *
Bystander administered naloxone	39 10%	<5 *
History of prior overdose	96 26%	16 21%

<5 Estimates with a count of less than 5 are suppressed due to statistical reliability and confidentiality purposes.

* Percentages are suppressed due to small cell sizes (n <5).

Key Drug Categories of Fatal Overdoses

Table 10 displays the frequencies of the most prominent drug categories causing death among confirmed drug deaths. Similarly to most other frequency distributions in this report, the drug category estimates are presented for the whole year of 2025 and for the rolling aggregate of the period January–March 2026. As expected, within the confirmed drug death cases in January–March 2026, nonpharmaceutical fentanyl, including nonpharmaceutical fentanyl analogs, was the most frequent cause of death, listed on the death certificate of 43 (58%) victims. The proportion of confirmed fentanyl overdose deaths has declined from its 2023 peak of 78% to 69% in 2024, and an estimated 54% so far for confirmed overdoses in 2025.

Fentanyl is nearly always found in combination with multiple other drugs. Xylazine was identified as a co-intoxicant with fentanyl for the first time in 2021. Among 393 confirmed deaths in 2025, there were 30 cases (8%) with xylazine listed in addition to fentanyl as a cause of death. During January–March 2026, there were fewer than 5 confirmed cases that involved xylazine in addition to fentanyl. In 2025, 2% of the confirmed drug deaths involved heroin. There was no mention of heroin involvement as a cause of death among confirmed cases in the last three months.

Stimulants continue to increase as a cause of death, usually in combination with other drugs, particularly fentanyl. Cocaine-involved fatalities constituted 36 (49%) of confirmed cases so far in January–March 2026. The percentage of cocaine cases among confirmed deaths in 2025 was 45%, which is higher than the percentage in 2024 (43%) and in 2023 (37%). Fentanyl is listed as a cause in combination with cocaine in 24 of the January–March 2026 cases, 67% of the cocaine cases. Methamphetamine was cited as a cause of death in 20 (27%) of the confirmed fatal overdoses in January–March 2026; 14 (70%) of the methamphetamine deaths also involved fentanyl as a co-intoxicant cause of death. Cocaine and methamphetamine are named together on 7 (9%) death certificates in the most recent three months.

Table 10: Key drug categories and combinations causing death among confirmed overdoses

Cause of death (alone or in combination with other drugs) Sample size for confirmed cases only	Jan–Dec 2025 Est. N = 393	Jan–Mar 2026 Est. N= 74
Fentanyl or fentanyl analogs	213 54%	43 58%
Heroin	8 2%	<5 *
Cocaine	178 45%	36 49%
Methamphetamine	122 31%	20 27%
Pharmaceutical opioids	90 23%	20 27%
Fentanyl and heroin	8 2%	<5 *
Fentanyl and cocaine	109 28%	24 32%
Fentanyl and methamphetamine	79 20%	14 19%
Fentanyl and xylazine	30 8%	<5 *
Cocaine and methamphetamine	41 10%	7 9%
Fentanyl, cocaine and methamphetamine	30 8%	5 7%

Note: Nonpharmaceutical tramadol is now being combined with fentanyl in pills and powders for illicit drug use. When found in combination with fentanyl, and in the absence of a known prescription, tramadol is categorized as a nonpharmaceutical opioid.

Emerging Substances Causing Death

The Maine Office of Chief Medical Examiner keeps close tabs on emerging substances that may cause death, most found in combination with fentanyl. When new substances are detected in Maine decedents by the National Medical Services Lab, the Office receives a report. Those substances are then quantified in the victim’s toxicology report and may be added as a cause of death. In most cases, Maine has not seen many of the new and emerging substances reported nationally. Frequencies have been very low, generally under 5. One exception is bromazolam, a nonpharmaceutical benzodiazepine, which was found in 10 (3%) of cases during 2025 and three cases so far in 2026. The other exception is ortho-methylfentanyl, a nonpharmaceutical fentanyl analog (an opioid), which was found in 8 (2%) of cases during 2025 and one case so far in 2026. Carfentanil (an opioid) is not new, but is a particularly lethal veterinary tranquilizer. It was found in 9 (2%) of 2025 Maine cases and two so far in 2026. Another dangerous veterinary sedative, medetomidine, which is not an opioid, is being found more often in other states, but was only found in one case during 2025 and none so far in 2026. Bromazolam and medetomidine overdoses cannot be reversed by naloxone.

Background Information about this Report

This report, funded jointly by the Maine Office of Attorney General and the Office of Behavioral Health,¹ provides an overview of statistics regarding suspected and confirmed fatal and nonfatal drug overdoses each month. Data for the fatal overdoses were collected at the Office of Chief Medical Examiner and data regarding nonfatal overdoses were contributed by the Maine CDC, Maine Emergency Medical Services, Maine ODMAP initiative, Maine Naloxone Distribution Initiative, and Office of Attorney General Naloxone Distribution. Year-to-date numbers are updated as medical examiner cases are finalized, and their overdose status is confirmed or ruled out, and as occasional lagged EMS, ED, and ODMAP data totals are finalized. The totals are expected to shift as case completion occurs. In addition, due to the small sample size in each month, we expect totals to fluctuate from month to month because of random variation. The monthly reports are posted on mainedrugdata.org.

A “drug death” is confirmed when one or more drugs are mentioned on the death certificate as a cause or significant contributing factor for the death. Most drug-induced fatalities are accidents related primarily to drug lethality, the unique vulnerability of the drug user, such as underlying medical conditions, and the circumstances surrounding drug use during that moment.

A “suspected” drug fatality is identified by physiological signs of overdose as well as physical signs at the scene and witness information. To be confirmed as a drug death, the medical examiner must have issued a final death certificate which includes the names of the specific drugs. A forensic toxicology exam must also have been done, which includes a minimum of two toxicology tests, one to screen for drugs present, and another that will quantify the levels of drugs in the decedent’s system. All cases receive a thorough external examination and comprehensive toxicology tests. In some cases, a complete autopsy is also done. Additional data, such as medical records and police incident reports are also collected. Normally cases are completed within one month; however, due to recent problems being experienced by our national toxicology testing service, completion of cases is occurring at about 6–8 weeks after death, and occasionally longer.

By highlighting drug deaths at the monthly level, this report brings attention to the often-dramatic shifts in totals that can occur from month to month. These fluctuations are common with small numbers and will tend toward an average over time. Starting in 2026, in order to support greater statistical reliability and due to confidentiality concerns, we will be using rolling three-month aggregations instead of single month totals in the tables. In addition, we will suppress totals less than 5.

Whereas the overall number of overdose deaths are a critical indicator of individual and societal stress, this metric itself can be quite resistant to public policy interventions due to its complexity. Overdose fatalities occur because of multiple unique and interacting factors, as mentioned above. For that reason, these reports will seek to monitor components that can be directly affected by specific public health education and harm reduction interventions. The statistics in this report reflect both suspected and confirmed “occurrent” deaths, that is, deaths that occur in the State of Maine, even though they may not be Maine residents. These totals also do not include Maine residents who die in other states. For these reasons, totals will differ slightly from the statistics reported by the National Center for Health Statistics, which reports only confirmed “resident” deaths. In addition, due to recently reported updates of toxicology results and newly confirmed or eliminated drug death cases, the 2025 statistics have changed slightly from those reported in the previous monthly report.

1 The Office of Attorney General supports ongoing research on fatal overdoses by the University of Maine. Additionally, the Overdose Data to Action cooperative agreement from the U.S. Centers for Disease Control & Prevention also provides funding to the State of Maine’s Office of Behavioral Health and Maine Center for Disease Control, which also supports University programs involving fatal and nonfatal overdoses surveillance and enables the collection of some of the nonfatal metrics included in this report. The conclusions in this report do not necessarily represent those of the U.S. Centers for Disease Control and Prevention.